

Microarrays in primary breast cancer – lessons from chemotherapy studies

P E Lønning, T Sørli^{1,2}, C M Perou², P O Brown³, D Botstein² and A-L Børresen-Dale¹

Section of Oncology, Department of Medicine, Haukeland University Hospital, Bergen, Norway

¹Department of Genetics, The Norwegian Radiumhospital, Oslo, Norway

²Department of Genetics, Stanford University School of Medicine, San Francisco, California, USA

³Department of Biochemistry and Howard Hughes Medical Institute, Stanford University School of Medicine, San Francisco, California, USA.

(C M Perou is now at Department of Genetics and Lineberger Comprehensive Cancer Center, The University of North Carolina, Chapel Hill, North Carolina, USA)

(Requests for offprints should be addressed to P E Lønning; Email: plon@haukeland.no)

Abstract

Current development in molecular techniques has extended the opportunities to explore genetic alterations in malignant tissue. There is a need to improve prognostication and, in particular, to understand the mechanisms of treatment resistance in different tumours. Gene analyses by microarrays allow concomitant analyses of several genes in concert, providing new opportunities for tumour classification and understanding of key biological disturbances. This paper outlines our continuing studies exploring prognostic and, we hope, predictive factors in breast cancer therapy.

Endocrine-Related Cancer (2001) 8 259–263

Introduction

Numerous prognostic factors have been studied in relation to breast cancer, but the biological knowledge that may be extracted from individual parameters is limited. First, many of these factors act in concert. Thus, multivariate analyses incorporating several factors generally reveal the prognostic impact of only two or three parameters, because of co-expression of individual parameters (Battaglia *et al.* 1988, Berger *et al.* 1988, Fisher *et al.* 1988, Fisher *et al.* 1993, Howat *et al.* 1983, Søreide *et al.* 1992). Secondly, because of this co-variance, prognostic factors do not necessarily reflect biological function. Thus, expressions of the oestrogen and the progesterone receptors are associated with a good prognosis in breast cancer, whether or not the patient receives adjuvant endocrine therapy (Harvey *et al.* 1999, Vollenweider-Zerargui *et al.* 1986). From a theoretical point of view this is unexpected, taking into account that ligand stimulation of the oestrogen receptor by oestradiol creates a potent mitogenic signal in cancer cells. Accordingly, the prognostic impact of oestrogen receptor expression is probably the result of its correlation to other parameters, such as low histological grade, slow growth rate (Singh *et al.* 1988), or other parameters.

There are several ways in which we might improve the way we are studying the biology of cancers *in vivo*.

Prognostic factors are limited by the fact that they often contain a 'predictive element'. The most important prognostic factor in breast cancer – expression of lymph node metastases – does not seem to be predictive of resistance or sensitivity to treatment (the reduction in the hazard ratio obtained with adjuvant chemotherapy or endocrine therapy seems to be approximately the same in lymph-node-positive and lymph-node-negative patients (Abe *et al.* 1998, Clarke *et al.* 1998). However, this is not the case for other parameters such as expression of the oestrogen receptor (Clarke *et al.* 1998) or mutations in the *TP53* gene (Geisler *et al.* 2001). From a therapeutic prospective, the major goal is to identify those patients whose tumours are sensitive and thus may benefit from treatment. An alternative to studying individual parameters is to look for several genetic alterations in concert, to better understand the functional network between individual parameters. This paper briefly views our current experience in applying such techniques to primary locally advanced breast cancer treated with neoadjuvant chemotherapy.

Clinical procedure

In 1991, we implemented our first procedure in which patients with locally advanced breast cancer (T₃/T₄, or N₂




Figure 1 Two-colour fluorescent hybridization for the analysis of gene expression.

tumours, or both) were treated with primary chemotherapy followed by local treatment (Aas *et al.* 1996). The bulk of these patients were senior patients with a median age of 64 years. Accordingly, we applied the drug regimen commonly used for metastatic breast cancer in our department at that time – namely, weekly doxorubicin monotherapy (dose 14 mg/m²). The scientific aim of this study was to explore biological markers in relation to chemoresistance. Accordingly, we preferred using a monotherapy regimen, in order to correlate these parameters to the effects of a particular drug.

Importantly, in this study we collected tumour samples before therapy by incisional biopsies and after chemotherapy (aimed at 16 weeks in accordance with the treatment procedure) when the patient underwent final surgery. In each case, tumours were collected and immediately snap-frozen in the operating theatre. Samples were stored in liquid nitrogen until required for processing.

Microarray techniques and analysis

The methods used have been outlined elsewhere (Alizadeh *et al.* 2000, DeRisi *et al.* 1997, Eisen & Brown 1999, Ross *et al.* 2000). cDNA prepared from mRNA in the experimental samples were labelled with Cy5, while the reference standard was made from a pool of mRNA isolated from 11 different cultured cell lines and labelled with Cy3 and the mixture was hybridised to microarrays with 8102 genes (Fig. 1). The microarray analysis was performed at the Stanford University Department of Genetics (P O Brown and D Botstein).

Results and discussion

In a previous study using this material (Aas *et al.* 1996), we showed that particular mutations in the *TP53* gene affecting the DNA-binding domains L2 and L3 predicted resistance to doxorubicin monotherapy. In a later extension of the study

A significant observation pertaining to the similarity in gene expression was that the 'paired samples' collected from the same tumour revealed a remarkable reproducibility with respect to gene profiling. Despite the fact that these samples were collected randomly from different parts of the tumour, that sample collection was separated by a time interval of 16 weeks, and that the patient had received chemotherapy in between, there was a remarkable consistency in gene expression between the two samples. Thus, when the tumours were classified by hierarchical clustering using the 1753 gene list, in 15 of the 20 pairs the two samples from the same tumour clustered next to each other, revealing a greater degree of similarity of gene expression (Fig. 2). Regarding the five pairs of samples for which such a similarity was not observed, in three cases this was characterised by a more 'normal-like' gene expression in the second sample. Notably, these three tumours were all collected from responders, meaning that the amount of tumour tissue compared with normal tissue could be significantly reduced in the second sample, which may explain this observation.

Continuing work and aims for future studies

Currently, we have extended the number of samples and are in the process of correlating profiles of gene expression to clinical outcome. A current observation is that the luminal group of tumours may be sub-divided into at least two sub-categories (luminal type A and B) with different gene expression profiles. By selecting tumours from our series of locally advanced cancers receiving uniform therapy for survival analysis, we were able to correlate tumour classification to clinical outcome (relapse-free and overall survival). As expected, the three oestrogen receptor-negative classes (basal-like, erbB-2-like and normal-cell-like tumours) were all associated with a poor outcome. Most interestingly, the two luminal sub-classes exhibited a significant difference in outcome with respect to relapse-free and overall survival, suggesting that this sub-classification may have novel clinical implications (Sørliie et al.).

In addition to the materials mentioned above, we have further analysed samples from another chemotherapy study of primary breast cancers, using a combined regimen of 5-fluorouracil and mitomycin, mainly in senior patients. Currently, we are correlating gene expression profiles in both series to clinical outcome, in particular with respect to drug responsiveness. We (Geisler et al. 2001) and others (Paik et al. 1998, Kandioler-Eckersberger et al. 2000) have shown that mutations in the *TP53* gene, and c-erbB-2 expression, correlate to resistance to chemotherapy in patients with breast cancer, but, importantly, we also observed tumours that expressed primary resistance to doxorubicin therapy despite harbouring wild type *TP53*. Most important, we also observed patients harbouring *TP53* mutations affecting the

DNA-binding domain who nevertheless responded to therapy (Geisler et al. 2001). Our current hypothesis is that, among patients expressing primary chemoresistance despite wild-type *TP53*, other disturbances in the p53 pathway may account for this phenotype. In addition, the finding that *TP53* mutations may be compensated for suggests that redundant mechanisms may be involved. Thus, a major aim of our current study programme is to evaluate other alterations in chemoresistance, in addition to *TP53* mutations. Clearly, drug resistance *in vivo* is a complex process likely to involve several genetic alterations and perhaps this also involves the patient's own genetic makeup. Accordingly, we hope that microarray studies of expression of several genes, together with complementary gene sequencing, could be a valuable tool for exploring this complex diversity.

Acknowledgements

The studies referred to were supported by the Norwegian Cancer Society, the Howard Hughes Medical Institute and the NCI.

References

- Aas T, Børresen A-L, Geisler S, Smith-Sørensen B, Johnsen H, Varhaug JE, Akslen LA & Lønning PE 1996 Specific P53 mutations are associated with *de novo* resistance to doxorubicin in breast cancer patients. *Nature Medicine* **2** 811–814.
- Abe O, Abe R, Enomoto K, Kikuchi K, Koyama H, Nomura Y, Sakai K, Sugimachi K, Tominaga T, Uchino J et al. 1998 Polychemotherapy for early breast cancer: an overview of the randomised trials. *Lancet* **TP53352** 930–942.
- Alizadeh AA, Eisen MB, Davis RE, Ma C, Lossos IS, Rosenwald A, Boldrick JC, Sabet H, Tran T, Yu X et al. 2000 Distinct types of diffuse large B-cell lymphoma identified by gene expression profiling. *Nature* **TP53403** 503–511.
- Battaglia F, Scambia G, Rossi S, Panici PB, Bellantone R, Polizzi G, Querzoli P, Negrini R, Iacobelli S, Crucitti F & Mancuso S 1988 Epidermal growth factor receptor in human breast cancer: correlation with steroid hormone receptors and axillary lymph node involvement. *European Journal of Cancer and Clinical Oncology* **24** 1685–1690.
- Berger MS, Locher GW, Saurer S, Gullick WJ, Waterfield MD, Groner B & Hynes NE 1988 Correlation of c-erbB-2 gene amplification and protein expression in human breast carcinoma with nodal status and nuclear grading. *Cancer Research* **48** 1238–1243.
- Clarke M, Collins R, Davies C, Godwin J, Gray R & Peto R 1998 Tamoxifen for early breast cancer: an overview of the randomised trials. *Lancet* **351** 1451–1467.
- DeRisi JL, Iyer VR & Brown PO 1997 Exploring the metabolic and genetic control of gene expression on a genomic scale. *Science* **278** 680–686.
- Eisen MB & Brown PO 1999 DNA arrays for analysis of gene expression. *Methods in Enzymology* **303** 179–205.
- Fisher B, Redmond C, Fisher ER & Caplan R 1988 Relative worth of estrogen or progesterone receptor and pathologic characteristics of differentiation as indicators of prognosis in

- node negative breast cancer patients: findings from national surgical adjuvant breast and bowel project protocol B-06. *Journal of Clinical Oncology* **6** 1076–1087.
- Fisher ER, Anderson S, Redmond C & Fisher B 1993 Pathologic findings from the national surgical adjuvant breast project protocol B-06. *Cancer* **71** 2507–2514.
- Geisler S, Lønning PE, Aas T, Johnsen H, Fluge O, Haugen DF, Lillehaug JR, Akslen LA & Borresen Dale AL 2001 Influence of TP53 gene alterations and c-erbB-2 expression on the response to treatment with doxorubicin in locally advanced breast cancer. *Cancer Research* **61** 2505–2512.
- Harvey JM, Clark GM, Osborne CK & Allred DC 1999 Estrogen receptor status by immunohistochemistry is superior to the ligand-binding assay for predicting response to adjuvant endocrine therapy in breast cancer. *Journal of Clinical Oncology* **17** 1474–1481.
- Howat JMT, Barnes DM, Harris M & Swindell R 1983 The association of cytosol oestrogen and progesterone receptors with histological features of breast cancer and early recurrence of disease. *British Journal of Cancer* **47** 629–640.
- Kandioler-Eckersberger D, Ludwig C, Rudas M, Kappel S, Janschek E, Wenzel C, Schlagbauer-Wadl H, Mittlböck M, Gnant M, Steger G & Jakesz R 2000 TP53 mutation and p53 overexpression for prediction of response to neoadjuvant treatment in breast cancer patients. *Clinical Cancer Research* **6** 50–56.
- Paik SM, Bryant J, Park CH, Fisher B, TanChiu E, Hyams D, Fisher ER, Lippman ME, Wickerham DL & Wolmark N 1998 erbB-2 and response to doxorubicin in patients with axillary lymph node-positive, hormone receptor-negative breast cancer. *Journal of the National Cancer Institute* **90** 1361–1370.
- Perou CM, Sørlie T, Eisen MB, van de Rijn M, Jeffrey SS, Rees CA, Pollack JR, Ross DT, Johnsen H, Akslen LA *et al.* 2000 Molecular portraits of human breast tumours. *Nature* **406** 747–752.
- Ross DT, Scherf U, Eisen MB, Perou CM, Rees C, Spellman P, Iyer V, Jeffrey SS, van de Rijn M, Waltham M *et al.* 2000 Systematic variation in gene expression patterns in human cancer cell lines. *Nature Genetics* **24** 227–235.
- Singh L, Wilson AJ, Baum M, Whimster WF, Birch IH, Jackson IM, Lowrey C & Palmer MK 1988 The relationship between histological grade, oestrogen receptor status, events and survival at 8 years in the NATO ('Novaldex') trial. *British Journal of Cancer* **57** 612–614.
- Søreide JA, Lea OA, Varhung JE, Skarstein A & Kvinnsland S 1992 Androgen receptors in operable breast cancer: relation to other steroid hormone receptors, correlations to prognostic factors and predictive value for effect of adjuvant tamoxifen treatment. *European Journal of Surgical Oncology* **18** 112–118.
- Sørlie T, Perou CK, Tibshirani R, Aas T, Geisler S, Johnsen H, Hastie T, Eisen MB, van den Rijn M, Jeffrey SS, Thorsen T, Quist H, Matese JC, Brown PO, Botstein D, Lønning PE & Borresen-Dale A-L 2001 Gene expression patterns of breast carcinomas distinguish tumor subclasses with clinical implications. *PNAS* (In Press).
- Vollenweider-Zerargui L, Barrelet L, Wong Y, Lemarchand-Béraud T & Gómez F 1986 The predictive value of estrogen and progesterone receptors' concentrations on the clinical behavior of breast cancer in women. Clinical correlation on 547 patients. *Cancer* **57** 1171–1180.

